arXiv:0801.1686v1 [gr-qc] 10 Jan 2008

Constraining a bulk viscous matter-dominated cosmological model using SNe Ia, CMB and LSS

Arturo Avelino, U. Nucamendi and F. S. Guzmán

Instituto de Física y Matemáticas
Universidad Michoacana de San Nicolás de Hidalgo
Edificio C-3, A.P. 2-82, Ciudad Universitaria, CP. 58040
Morelia, Michoacán, México

Abstract

We present and constrain a cosmological model which component is a pressureless fluid with bulk viscosity as an explanation for the present accelerated expansion of the universe. We study the particular model of a constant bulk viscosity coefficient $\zeta_{\rm m}$. The possible values of $\zeta_{\rm m}$ are constrained using the cosmological tests of SNe Ia Gold 2006 sample, the CMB shift parameter R from the three-year WMAP observations, the Baryon Acoustic Oscillation (BAO) peak A from the Sloan Digital Sky Survey (SDSS) and the Second Law of Thermodynamics (SLT). It was found that this model is in agreement with the SLT using only the SNe Ia test. However when the model is submitted to the three cosmological tests together (SNe+CMB+BAO) the results are: 1.- the model violates the SLT, 2.- predicts a value of $H_0 \approx 53 \ {\rm km} \cdot {\rm sec}^{-1} \cdot {\rm Mpc}^{-1}$ for the Hubble constant, and 3.- we obtain a bad fit to data with a $\chi^2_{\rm min} \approx 400$ ($\chi^2_{\rm d.o.f.} \approx 2.2$). These results indicate that this model is ruled out by the observations.

We present a flat cosmological model which component is a pressureless fluid made of baryon and dark matter components with a constant bulk viscosity coefficient of the form $\zeta_{\rm m}$ =constant, which potentially could explain the present accelerated expansion of the universe. A bulk viscosity coefficient can produce a positive term in the second Friedmann equation that induces an acceleration [1, 2] (i.e., $\ddot{a} \geq 0$, with a denoting the scale factor and the dots derivatives with respect to the cosmic time). Similar models or analysis have been proposed also in [2, 3, 4, 5].

Model of the bulk viscous matter-dominated universe The energy-momentum tensor of a fluid composed by *only* matter (hereinafter, we call *matter* to baryon and dark matter components together) with bulk viscosity $\zeta_{\rm m}$ is defined by $T_{\mu\nu} = \rho_{\rm m} u_\mu u_\nu + (g_{\mu\nu} + u_\mu u_\nu) P^*_{\rm m}$, where $P^*_{\rm m} = P_{\rm m} - 3\zeta_{\rm m} \, (\dot{a}/a)$ (see e.g. [1]). Here $P_{\rm m}$ is the pressure of the matter fluid. However, assuming pressureless matter: $P_{\rm m} = 0 \ \Rightarrow \ P^*_{\rm m} = -3\zeta_{\rm m} \, (\dot{a}/a)$. The subscript "m" stands for *matter.* The four-velocity vector u_μ is that of a comoving observer that measures the pressure $P^*_{\rm m}$ and the density $\rho_{\rm m}$ of the matter fluid. The conservation of energy has the form $a(d\rho_{\rm m}/da) = 3\,(3\zeta_{\rm m} H - \rho_{\rm m})$, where $H \equiv \dot{a}/a$ is the Hubble parameter. This equation in terms of the redshift reads $(1+z)\left(d\hat{\Omega}_{\rm m}/dz\right) + \hat{\Omega}_{\rm m}^{1/2}\;\tilde{\zeta}_0 - 3\hat{\Omega}_{\rm m} = 0$, where we have defined [1] $\hat{\Omega}_{\rm m}(z) \equiv \rho_{\rm m}/\rho^0_{\rm crit}$, and the dimensionless coefficient $\tilde{\zeta}_0 \equiv \zeta_{\rm m}(24\pi G/H_0)$. The exact solution to this equation is $\hat{\Omega}_{\rm m}(z) = (1/9)\left[(\tilde{\zeta}_0 - 3)(1+z)^{3/2} - \tilde{\zeta}_0\right]^2$. On the other hand, a spatially flat geometry is assumed ($k = 0$) for the FRW metric cosmology as suggested by WMAP. Thus, the first Friedmann equation for a flat universe composed by bulk viscous matter has the form: $H^2(z) = (8\pi G/3)\rho_{\rm m} = H_0^2\,\hat{\Omega}_{\rm m}(z)$. Therefore $H(z) = (H_0/3)\left[\left(\tilde{\zeta}_0 - 3\right)(1+z)^{3/2} - \tilde{\zeta}_0\right]$.

Cosmological tests We constrain the possible values of the bulk viscosity $\tilde{\zeta}_0$ using the cosmological tests of the Gold 2006 SNe Ia data sample [6] composed by 182 SNe Ia, the Cosmic Microwave Background (CMB) shift parameter R from the three-year WMAP observations, the Baryon Acoustic Oscillation (BAO) peak A from the Sloan Digital Sky Survey (SDSS) and the Second Law of Thermodynamics (SLT).

For the SNe Ia we define the observational luminosity distance [7, 8] in a flat cosmology and with the bulk viscous matter component as $d_L = c(1+z)H_0^{-1}\int_0^z E(z')^{-1}\,dz'$, where $E(z) \equiv H(z)/H_0$. The *theoretical distance moduli* for the i-th supernovae with redshift z_i is $\mu^{\rm t}(z_i) = 5\log_{10}[d_L(z_i)/1{\rm Mpc}] + 25$, with c the speed of light. The $\chi^2_{\rm SNe}$ statistical function becomes $\chi^2_{\rm SNe}(\tilde{\zeta}_0, H_0) \equiv \sum_{k=1}^{182}\left[\mu^{\rm t}(z_k, \tilde{\zeta}_0, H_0) - \mu_k\right]^2/\sigma_k^2$, where μ_k is the *observed* distance moduli for the k-th supernovae and σ_k^2 is the variance of the measurement.

The CMB shift parameter R is defined as $R \equiv \sqrt{\Omega^0_{\rm m}}\int_0^{Z_{\rm CMB}} E(z')^{-1}\,dz'$, where $Z_{\rm CMB} = 1089$ is the redshift of recombination [9, 10]. The observed value of the shift parameter R is reported to be $R_{\rm obs} = 1.70 \pm 0.03$ [10].

From the LSS data, we use the baryon acoustic oscillation (BAO) peak A defined as $A \equiv \sqrt{\Omega^0_{\rm m}}E(z_1)^{-1/3}\left[z_1^{-1}\int_0^{z_1} E(z')^{-1}\,dz'\right]^{2/3}$, where $z_1 = 0.35$ [11]. The observed value of SDSS-BAO peak is $A_{\rm obs} = 0.469 \pm 0.017$ [11].

We construct the joint χ^2 function as $\chi^2_{\rm total} = \chi^2_{\rm SNe} + \chi^2_{\rm CMB} + \chi^2_{\rm BAO}$, where $\chi^2_{\rm SNe}$ was defined above, and $\chi^2_{\rm CMB} = [(R - R_{\rm obs})/\sigma_R]^2$ and $\chi^2_{\rm BAO} = [(A - A_{\rm obs})/\sigma_A]^2$.

The generation of *local* entropy in the space–time is defined as $T\,\nabla_\mu s^\mu = \zeta\nabla_\mu u^\mu = 3H\zeta$ [1], where T is the temperature, $\nabla_\mu s^\mu$ is the rate at which entropy is being generated in a unit volume and ζ is the bulk viscosity coefficient. The second law of thermodynamics can be written as $T\nabla_\mu s^\mu \geq 0 \;\Rightarrow\; 3H\zeta \geq 0$. Since H is positive in an expanding universe then ζ has to be positive in order to

---

[1] We use this particular definition of $\hat{\Omega}_{\rm m}$ for computational conveniences. $\rho^0_{\rm critic} \equiv 3H_0^2/8\pi G$ is the *critical density* today, where H_0 is the Hubble constant.



Figure 1: Joint confidence intervals for the parameters $(\tilde{\zeta}_0, H_0)$ for a bulk viscous matter-dominated universe, spatially flat and without any dark energy component. We show the confidence intervals of $68.3\%(1\sigma)$, $95.4\%(3\sigma)$ and $99.73\%(5\sigma)$. The constraints are derived from the Gold 2006 SNe Ia data sample alone (left), and the joint Gold 2006 + CMB + BAO cosmological tests (right).

preserve the validity of the second law of thermodynamics. Thus, in the present model a necessary condition is $\zeta_{\rm m} \geq 0$.

Constraining the bulk viscous cosmological model We compute the *best estimated values* and "the goodness-of-fit" of $\tilde{\zeta}_0$ and H_0 to the data through χ^2-minimization, using SNe alone, and also considering the CMB and BAO. Then we compute the confidence intervals for $(\tilde{\zeta}_0, H_0)$ to constrain their possible values. We obtain as best estimates using only SNe Ia: $(\tilde{\zeta}_0 = 1.71, H_0 = 62.30)$ with a $\chi^2_{\rm min} = 161.2$ $(\chi^2_{\rm d.o.f.} = 0.895)$, where hereinafter H_0 is in units of $\,\rm km \cdot sec^{-1} \cdot Mpc^{-1}$. And using SNe Ia, CMB and BAO together: $(\tilde{\zeta}_0 = -0.30, H_0 = 52.97)$ with a $\chi^2_{\rm min} = 400.5$ $(\chi^2_{\rm d.o.f.} = 2.225)$.

It can be seen that when we consider only SNe Ia we obtain $\tilde{\zeta}_0 \geq 0$ with a 99.7% confidence level and a reasonable value for the Hubble constant in the interval $59.5 \geq H_0 \geq 65.4$ as well as for $\chi^2_{\rm d.o.f.} = 0.895$. However, when the same analysis using the three cosmological tests together (SNe Ia, CMB and BAO) is performed, we obtain negative values of $\tilde{\zeta}_0$ with at least 99.7 % confidence level that disagree with the SLT, a not so good estimation for H_0 of $51.4 \geq H_0 \geq 54.6$, and a bad $\chi^2_{\rm d.o.f.} = 2.225$. These results are illustrated in Figure 1.

Therefore, it is possible to conclude that this model is not a good candidate for explaining the observed accelerated expansion of the universe as suggested in [2, 4, 5].

Acknowledgments This work is partly supported by grants CIC-UMSNH 4.8, 4.9, PROMEP UMSNH-CA-22, SNI-20733 and COECYT CB0702137-2.

References

[1] C. W. Misner, K. S. Thorne and J. A. Wheeler, **Gravitation**, W. H. Freeman and Company, p 567, (1973).

[2] J. C. Fabris, S. V. B. Goncalves and R. de Sa Ribeiro, *Gen. Rel. Grav* **38**, 495–506, (2006).

[3] Arturo Avelino and Ulises Nucamendi, *in preparation*, (2008).

[4] R. Colistete et al., *Phys. Rev. D* **76**, 103516, 1–13, (2007).

[5] Jie Ren and Xin-He Meng, *Phys. Lett. B* **633**, 1–8, (2006).

[6] Adam G. Riess et al., `arXiv:astro-ph/0611572v2`, (2006).

[7] M. Turner and A. G. Riess, *Astrophysical journal* **569**, 18, (2002).

[8] A. G. Riess et al., *Astrophysical journal* **607**, 665–687, (2004)

[9] A. Melchiorri et al., *Phys. Rev. D* **68**, 043509, (2003).

[10] Y. Wang and P. Mukherjee, *Astrophysical Journal* **650**, 1–6, (2006).

[11] D. J. Eisenstein et al., (SDSS Collaboration) *Astrophysical Journal* **633**, 560–574, (2005).